

T.E.D.[™]Anti-Embolism Stockings DVT Prevention for your patient

How to Measure

Studies that proved the clinical effectiveness of T.E.D.[™] anti-embolism stockings used thigh length styles. Select thigh length wherever possible unless knee length is more appropriate to patient condition.



1 Measure Upper Thigh Circumference

If thigh circumference is less than 91.4cm, choose: Thigh length or Thigh length with belt

2 Measure Calf Circumference at greatest point3 Measure distance from Buttock Fold to Heel

If Upper Thigh Circumference is 91.4cm or more

choose: Knee length

2 Measure Calf Circumference at greatest point

3 Measure distance from behind Knee to Heel

Selection Guide

Thigh length Style Thigh Circumfe eg Length 3Calf Girth 2Less than 74cm Short 3071LF A Less than 74cm to 84cm Regular 3130LF 30.5cm В Small 3222LF 84cm or more Long С Less than 74cm Short 3310LF D 30.5 to 38cm Less than 63.5cm 74cm to 84cm Regular 3416LF E Medium 84cm or more Long 3549LF E 3634LF Less than 74cm Short G 38 to 44.5cm 74cm to 84cm Regular 3728LF H Large 3856LF 84cm or more Long J Less than 74cm Short 4010LF K 38 to 44.5cm 63.5 to 81.3cm 74cm to 84cm Regular 4114LF L Extra Large 84cm or more Long 4216LF M 3180LF Ν Less than 74cm Short 44.5 to 54.6cm 3181LF 63.5 to 81.3cm 74cm to 84cm Regular Ρ Extra Large Plus 84cm or more Long 3182LF Q Less than 74cm Short 3183LF R 54.6 to 66cm 81.3 to 91.4cm 74cm to 84cm Regular 3184LF S Extra Extra Large 3185LF 84cm or more Long Т

IMPORTANT - to ensure your patient gets the right levels of compression, it is essential to take accurate measurements. By following the measuring guide you will be able to select the correct style and size of stocking.

Fitting Guide



Put hand inside stocking and grab heel pocket.



Pull stocking inside out.



Thigh length with Belt Style							
Thigh Circumference 1	Calf Girth 2	Leg Length 3	Code	Size	Colou Toe	Colour Toe Top	
Less than 63.5cm	Less than 25cm Extra Small	Less than 71cm Regular	3306	-	100	Top	
		71cm & more Long	3320	-			
	25 to 30.5cm Small	Less than 72cm Regular	3039-	A+			
		72cm & more Long	3364	B+			
	30.5 to 38cm	Less than 72cm Regular	3144	C+			

72 cm & more Long

Medium




Place stocking over foot. Align inspection toe hole to fall under toes. Ensure patient's heel is centred in the heel pocket.

After calf portion is applied start rotating stocking inward where the break in pressure occurs, so that the gusset is placed over the femoral artery (slightly towards the inside of the thigh). The stitch change in the thigh length should be positioned 3-5cm below knee (for knee length style, position top 3-5cm below knee).

Download the **FREE T.E.D.[™] APP** and discover sizing and application procedures.





		72cm & more Long	J449	D+	
	38 to 44.5cm	Less than 74cm Regular	3221	E+	
	Large	74 cm & more Long	3523	F+	
63.5 to 81.3cm	38 to 44.5cm Extra Large	Less than 72cm Regular	3922	G+	
05.5 10 81.5011		72cm & more Long	3995	H+	

3449 D+

Stockings should be laundered every 2 to 3 days, unless soiled (in case they should be cleaned/replaced immediately). Laundering increases the length of service by removing body secretions from the elastic threads.

Knee length Style						
Calf Girth 2	Leg Length 3	Code	Size	Colour		
	Leg Length 🥥			Тое	Тор	
Less than 30.5cm Small	Less than 41cm Regular	7071-	A-			
	41cm & more Extra Long	7339	B-			
30.5 to 38cm Medium	Less than 43cm Regular	7115	C-			
	43cm & more Extra Long	7480	D-			
38 to 44.5cm Large	Less than 46cm Regular	7203-	E-			
	46cm & more Extra Long	7594	F-			
44.5 to 51cm Extra Large	Less than 46cm Regular	7604-	G-			
	46cm & more Extra Long	7802-	H-			
51 to 58.4cm Extra Extra Large	Less than 46cm Regular	7470LF	J-			
	46cm & more Extra Long	7471LF	K-			
58.4 to 66cm Extra Extra Extra Large	Less than 46cm Regular	7472LF	L-			
	46cm & more Extra Long	7473LF	M-			

CONTRA-INDICATIONS: Massive oedema of leg or pulmonary oedema from congestive heart failure; severe arteriosclerosis or other ischaemic vascular disease; extreme deformity of the leg. Any local leg conditions in which stockings would interfere, such as:

1) Dermatitis3) Gangrene2) Vein Ligation (immediate postoperative)4) Recent Skin Graft